

2022 AMC 12A

For more practice and resources, visit ziml.areteem.org

The problems in the AMC-Series Contests are copyrighted by American Mathematics Competitions at Mathematical Association of America (www.maa.org).



Question 1

Not yet answered

Points out of 6

What is the value of

$$3+\frac{1}{3+\frac{1}{3+\frac{1}{3}}}?$$

$\textbf{(A)}\ \frac{31}{10}$ $\textbf{(B)}\ \frac{49}{15}$ $\textbf{(C)}\ \frac{33}{10}$ $\textbf{(D)}\ \frac{109}{33}$ $\textbf{(E)}\ \frac{15}{4}$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 2

Not yet answered

Points out of 6

The sum of three numbers is 96. The first number is 6 times the third number, and the third number is 40 less than the second number. What is the absolute value of the difference between the first and second numbers?

$\textbf{(A)}\ 1$ $\textbf{(B)}\ 2$ $\textbf{(C)}\ 3$ $\textbf{(D)}\ 4$ $\textbf{(E)}\ 5$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 3

Not yet answered

Points out of 6

Five rectangles, A, B, C, D, and E, are arranged in a square as shown below. These rectangles have dimensions 1×6, 2×4, 5×6, 2×7, and 2×3, respectively. (The figure is not drawn to scale.) Which of the five rectangles is the shaded one in the middle?



$\textbf{(A)}\ A \qquad \textbf{(B)}\ B \qquad \textbf{(C)}\ C \qquad \textbf{(D)}\ D \qquad \textbf{(E)}\ E$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 4

Not yet answered

Points out of 6

The least common multiple of a positive divisor n and 18 is 180, and the greatest common divisor of n and 45 is 15. What is the sum of the digits of n?

$\textbf{(A)}\ 3 \qquad \textbf{(B)}\ 6 \qquad \textbf{(C)}\ 8 \qquad \textbf{(D)}\ 9 \qquad \textbf{(E)}\ 12$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 5

Not yet answered

Points out of 6

The $taxicab\ distance$ between points (x_1, y_1) and (x_2, y_2) in the coordinate plane is given by $|x_1 - x_2| + |y_1 - y_2|$. For how many points P with integer coordinates is the taxicab distance between P and the origin less than or equal to 20?

$\mathbf{(A)}\ 441 \qquad \mathbf{(B)}\ 761 \qquad \mathbf{(C)}\ 841 \qquad \mathbf{(D)}\ 921 \qquad \mathbf{(E)}\ 924$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 6

Not yet answered

Points out of 6

A data set consists of 6 (not distinct) positive integers: 1, 7, 5, 2, 5, and X. The average (arithmetic mean) of the 6 numbers equals a value in the data set. What is the sum of all positive values of X?

$\mathbf{(A)}\ 10 \qquad \mathbf{(B)}\ 26 \qquad \mathbf{(C)}\ 32 \qquad \mathbf{(D)}\ 36 \qquad \mathbf{(E)}\ 40$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 7

Not yet answered

Points out of 6

A rectangle is partitioned into 5 regions as shown. Each region is to be painted a solid color - red, orange, yellow, blue, or green - so that regions that touch are painted different colors, and colors can be used more than once. How many different colorings are possible?



$\textbf{(A)}\ 120 \qquad \textbf{(B)}\ 270 \qquad \textbf{(C)}\ 360 \qquad \textbf{(D)}\ 540 \qquad \textbf{(E)}\ 720$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 8

Not yet answered

Points out of 6

The infinite product

$$\sqrt[3]{10} \cdot \sqrt[3]{\sqrt[3]{10}} \cdot \sqrt[3]{\sqrt[3]{\sqrt[3]{10}}} \ldots$$

evaluates to a real number. What is that number?

$\textbf{(A)}\ \sqrt{10} \qquad \textbf{(B)}\ \sqrt[3]{100} \qquad \textbf{(C)}\ \sqrt[4]{1000} \qquad \textbf{(D)}\ 10 \qquad \textbf{(E)}\ 10\sqrt[3]{10}$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 9

Not yet answered

Points out of 6

On Halloween 31 children walked into the principal's office asking for candy. They can be classified into three types: Some always lie; some always tell the truth; and some alternately lie and tell the truth. The alternaters arbitrarily choose their first response, either a lie or the truth, but each subsequent statement has the opposite truth value from its predecessor. The principal asked everyone the same three questions in this order.

"Are you a truth-teller?" The principal gave a piece of candy to each of the 22 children who answered yes.

"Are you an alternater?" The principal gave a piece of candy to each of the 15 children who answered yes.

"Are you a liar?" The principal gave a piece of candy to each of the 9 children who answered yes.

How many pieces of candy in all did the principal give to the children who always tell the truth?

$\textbf{(A)}\ 7 \qquad \textbf{(B)}\ 12 \qquad \textbf{(C)}\ 21 \qquad \textbf{(D)}\ 27 \qquad \textbf{(E)}\ 31$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 10

Not yet answered

Points out of 6

How many ways are there to split the integers 1 through 14 into 7 pairs such that in each pair, the greater number is at least 2 times the lesser number?

$\textbf{(A)}\ 108 \qquad \textbf{(B)}\ 120 \qquad \textbf{(C)}\ 126 \qquad \textbf{(D)}\ 132 \qquad \textbf{(E)}\ 144$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 11

Not yet answered

Points out of 6

What is the product of all real numbers x such that the distance on the number line between $\log_6 x$ and $\log_6 9$ is twice the distance on the number line between $\log_6 10$ and 1?

$\textbf{(A)}\ 10 \qquad \textbf{(B)}\ 18 \qquad \textbf{(C)}\ 25 \qquad \textbf{(D)}\ 36 \qquad \textbf{(E)}\ 81$

Select one:

- A
- B
- C
- D
- E
- Leave blank (1.5 points)

Question 12

Not yet answered

Points out of 6

Let M be the midpoint of AB in regular tetrahedron $ABCD$. What is $\cos(\angle CMD)$?

$\textbf{(A)}\ \frac{1}{4} \qquad \textbf{(B)}\ \frac{1}{3} \qquad \textbf{(C)}\ \frac{2}{5} \qquad \textbf{(D)}\ \frac{1}{2} \qquad \textbf{(E)}\ \frac{\sqrt{3}}{2}$

Select one:

- A
- B
- C
- D
- E
- Leave blank (1.5 points)

Question 13

Not yet answered

Points out of 6

Let $\mathcal{R}$ be the region in the complex plane consisting of all complex numbers z that can be written as the sum of complex numbers z_1 and z_2, where z_1 lies on the segment with endpoints 3 and $4i$, and z_2 has magnitude at most 1. What integer is closest to the area of $\mathcal{R}$?

$\textbf{(A)}\ 13 \qquad \textbf{(B)}\ 14 \qquad \textbf{(C)}\ 15 \qquad \textbf{(D)}\ 16 \qquad \textbf{(E)}\ 17$

Select one:

- A
- B
- C
- D
- E
- Leave blank (1.5 points)

Question 14

Not yet answered

Points out of 6

What is the value of

$$(\log 5)^3 + (\log 20)^3 + (\log 8)(\log 0.25)$$

where $\log$ denotes the base-ten logarithm?

(A) $\frac{3}{2}$ **(B)** $\frac{7}{4}$ **(C)** 2 **(D)** $\frac{9}{4}$ **(E)** 3

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 15

Not yet answered

Points out of 6

The roots of the polynomial $10x^3 - 39x^2 + 29x - 6$ are the height, length, and width of a rectangular box (right rectangular prism). A new rectangular box is formed by lengthening each edge of the original box by 2 units. What is the volume of the new box?

(A) $\frac{24}{5}$ **(B)** $\frac{42}{5}$ **(C)** $\frac{81}{5}$ **(D)** 30 **(E)** 48

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 16

Not yet answered

Points out of 6

A triangular number is a positive integer that can be expressed in the form $t_n = 1 + 2 + 3 + \cdots + n$, for some positive integer n. The three smallest triangular numbers that are also perfect squares are $t_1 = 1 = 1^2$, $t_8 = 36 = 6^2$, and $t_{49} = 1225 = 35^2$. What is the sum of the digits of the fourth smallest triangular number that is also a perfect square?

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 17

Not yet answered

Points out of 6

Supppose a is a real number such that the equation

$$a \cdot (\sin x + \sin(2x)) = \sin(3x)$$

has more than one solution in the interval $(0, \pi)$. The set of all such a that can be written in the form

$$(p, q) \cup (q, r),$$

where p, q, and r are real numbers with $p < q < r$. What is $p + q + r$?

$\textbf{(A)}\ -4 \qquad \textbf{(B)}\ -1 \qquad \textbf{(C)}\ 0 \qquad \textbf{(D)}\ 1 \qquad \textbf{(E)}\ 4$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 18

Not yet answered

Points out of 6

Let T_k be the transformation of the coordinate plane that first rotates the plane k degrees counterclockwise around the origin and then reflects the plane across the y-axis. What is the least positive integer n such that performing the sequence of transformations $T_1, T_2, T_3, \cdots, T_n$ returns the point $(1, 0)$ back to itself?

$\textbf{(A)}\ 359 \qquad \textbf{(B)}\ 360 \qquad \textbf{(C)}\ 719 \qquad \textbf{(D)}\ 720 \qquad \textbf{(E)}\ 721$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 19

Not yet answered

Points out of 6

Suppose that 13 cards numbered $1, 2, 3, \ldots, 13$ are arranged in a row. The task is to pick them up in numerically increasing order, working repeatedly from left to right. In the example below, cards $1, 2, 3$ are picked up on the first pass, 4 and 5 on the second pass, 6 on the third pass, $7, 8, 9, 10$ on the fourth pass, and $11, 12, 13$ on the fifth pass. For how many of the $13!$ possible orderings of the cards will the 13 cards be picked up in exactly two passes?

7	11	8	6	4	5	9	12	1	13	10	2	3

$\textbf{(A)}\ 4082 \qquad \textbf{(B)}\ 4095 \qquad \textbf{(C)}\ 4096 \qquad \textbf{(D)}\ 8178 \qquad \textbf{(E)}\ 8191$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 20

Not yet answered

Points out of 6

Isosceles trapezoid $ABCD$ has parallel sides $\overline{AD}$ and $\overline{BC}$, with $BC < AD$ and $AB = CD$. There is a point P in the plane such that $PA = 1, PB = 2, PC = 3$, and $PD = 4$. What is $\frac{BC}{AD}$?

$\textbf{(A)}\ \frac{1}{4} \qquad \textbf{(B)}\ \frac{1}{3} \qquad \textbf{(C)}\ \frac{1}{2} \qquad \textbf{(D)}\ \frac{2}{3} \qquad \textbf{(E)}\ \frac{3}{4}$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 21

Not yet answered

Points out of 6

Let

$$P(x) = x^{2022} + x^{1011} + 1.$$

Which of the following polynomials is a factor of $P(x)$?

$\textbf{(A)}\ x^2 - x + 1 \qquad \textbf{(B)}\ x^2 + x + 1 \qquad \textbf{(C)}\ x^4 + 1$
$\textbf{(D)}\ x^6 - x^3 + 1 \qquad \textbf{(E)}\ x^6 + x^3 + 1$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 22

Not yet answered

Points out of 6

Let c be a real number, and let z_1 and z_2 be the two complex numbers satisfying the equation $z^2 - cz + 10 = 0$. Points z_1, z_2, $\frac{1}{z_1}$, and $\frac{1}{z_2}$ are the vertices of (convex) quadrilateral Q in the complex plane. When the area of Q obtains its maximum possible value, c is closest to which of the following?

$\textbf{(A)}\ 4.5 \qquad \textbf{(B)}\ 5 \qquad \textbf{(C)}\ 5.5 \qquad \textbf{(D)}\ 6 \qquad \textbf{(E)}\ 6.5$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 23

Not yet answered

Points out of 6

Let h_n and k_n be the unique relatively prime positive integers such that

$$\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} = \frac{h_n}{k_n}.$$

Let L_n denote the least common multiple of the numbers $1, 2, 3, \cdots, n$. For how many integers n with $1 \leq n \leq 22$ is $k_n < L_n$?

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 24

Not yet answered

Points out of 6

How many strings of length 5 formed from the digits 0, 1, 2, 3, 4 are there such that for each $j \in \{1, 2, 3, 4\}$, at least j of the digits are less than j? (For example, 02214 satisfies this condition because it contains at least 1 digit less than 1, at least 2 digits less than 2, at least 3 digits less than 3, and at least 4 digits less than 4. The string 23404 does not satisfy the condition because it does not contain at least 2 digits less than 2.)

$\textbf{(A)}\ 500 \qquad \textbf{(B)}\ 625 \qquad \textbf{(C)}\ 1089 \qquad \textbf{(D)}\ 1199 \qquad \textbf{(E)}\ 1296$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**

Question 25

Not yet answered

Points out of 6

A circle with integer radius r is centered at (r, r). Distinct line segments of length c_i connect points $(0, a_i)$ to $(b_i, 0)$ for $1 \leq i \leq 14$ and are tangent to the circle, where a_i, b_i, and c_i are all positive integers and $c_1 \leq c_2 \leq \cdots \leq c_{14}$. What is the ratio $\frac{c_{14}}{c_1}$ for the least possible value of r?

$\textbf{(A)}\ \frac{21}{5} \qquad \textbf{(B)}\ \frac{85}{13} \qquad \textbf{(C)}\ 7 \qquad \textbf{(D)}\ \frac{39}{5} \qquad \textbf{(E)}\ 17$

Select one:

- **A**
- **B**
- **C**
- **D**
- **E**
- **Leave blank (1.5 points)**